Saskatchewan
Ministry of
Agriculture

Crop Report

Published by the Ministry of Agriculture
ISSN 0701 7085
Report number 21, August 23, 2012

For the Period August 14 to 20, 2012

Warm and dry weather in many areas of the province has allowed producers to combine 11 per cent of the 2012 crop. Eighteen per cent is swathed or ready to straight combine, according to Saskatchewan Agriculture's weekly Crop Report.

The five-year (2007-2011) average for this time of year is eight per cent combined and 15 per cent swathed or ready to straight combine. Last year at this time, nine per cent had been combined and 18 per cent swathed or ready to straight combine.

Seventy-nine per cent of the winter wheat, 78 per cent of the fall rye, four per cent of the spring wheat, seven per cent of the durum, six per cent of the barley, five per cent of the canola, 22 per cent of the mustard, 36 per cent of the lentils and 41 per cent of the field peas have been combined. Thirty-two per cent of the canola, 26 per cent of the mustard, 15 per cent of the winter wheat, 12 per cent of the fall rye, 20 per cent of the lentils and 21 per cent of the field peas are swathed or ready to straight combine.

Across the province, topsoil moisture on cropland is rated as five per cent surplus, 72 per cent adequate, 19 per cent short and four per cent very short. Hay land and pasture topsoil moisture is rated as three per cent surplus, 64 per cent adequate, 24 per cent short and nine per cent very short.

Farmers are busy swathing, desiccating and combining and hauling bales.

One year ago

Nine per cent of the 2011 crop had been combined and 18 per cent was swathed or ready to straight combine. Warm and dry weather helped to progress harvest. Follow the 2012 Crop Report on Twitter @SKGovAg

Harvest Progress in SK Per cent combined	
All Crops	
Aug 23/12	11
5 year avg. (2007-2011)	8
Aug 25/11	9
Aug 23/10	2
Aug 25/09	3
Aug 24/08	9
Aug 19/07	16

Saskatchewan Crop Insurance Corporation Report – August 22

The deadline to select winterkill coverage for fall rye and winter wheat is August 25.

For further information, contact: Grant McLean, PAg,
Cropping Management Specialist, Agriculture Knowledge Centre,
Toll Free: 1-866-457-2377, E-mail: cropreport@gov.sk.ca.
Also available on the Ministry of Agriculture website at www.agriculture.gov.sk.ca.

Agriculture and Agri-Food Canada | Agriculture et Agroalimentaire Canada

SCIC
SASKATCHEWAN CROP INSURANCE CORPORATION

Southeastern Saskatchewan (Crop Districts 1, 2, and 3ASE)

Warm and dry weather this past week is moving harvest along. The region received very little precipitation; at 4 mm, the Moose Jaw area received the greatest amount of rain. The Odessa and Indian Head areas received 2 mm and the Wilcox area 3 mm. According to Crop Report precipitation information, the Tantallon area has the second highest cumulative rainfall in the province since April 1 (531 mm). In contrast, the Radville area has the lowest cumulative rainfall in the province (190 mm).

Harvest is quickly progressing and producers now have 20 per cent of the 2012 crop harvested, up from five per cent last week. Twenty-six per cent of the crop is swathed or ready to straight combine. Eighty-three per cent of the winter wheat, 78 per cent of the fall rye, 12 per cent of the spring wheat, 13 per cent of the durum, 10 per cent of the barley, 40 per cent of the lentils, 54 per cent of the field peas, five per cent of the chickpeas, 14 per cent of the canola and two per cent of the mustard have been combined. Fifty per cent of the canola and 46 per cent of the mustard has been swathed. Crop conditions and staging vary throughout the region, depending on the excess moisture in the spring, seeding date and the impact from disease.

Many crop reporters are indicating that it is difficult to assess yields at this time on uncombined crops due to the variable crop staging, disease and heat stress damage. Most crop yield estimates have decreased in the past few weeks. Average yields for the region are predicted as follows: winter wheat 54 bu/ac, spring wheat 35 bu/ac, durum 36 bu/ac, oats 64 bu/ac, barley 54 bu/ac, fall rye 47 bu/ac, flax 21 bu/ac, canola 28 bu/ac, mustard 840 lb./ac, sunflowers 1650 lb./ac, lentils 1250 lb./ac, peas 32 bu/ac, canaryseed 1014 lb./ac and chickpeas 2066 lb./ac.

Topsoil moisture ratings on cropland are reported as two per cent surplus, 80 per cent adequate and 18 per cent short. On hay land and pasture, topsoil moisture is rated as one per cent surplus, 76 per cent adequate and 23 per cent short. Crop District 3ASE is reporting that 35 per cent of cropland and 47 per cent of hay land and pasture are short moisture.

The majority of crop damage this week is due to wind, localized flooding, animals, insects and disease. Aster yellows, sclerotinia, fusarium, lodging and July's high temperatures will affect yields. Many producers in the region are indicating that yields on harvested crops are less than had been anticipated. There have been reports of grasshoppers in standing flax fields and wildlife feeding on crops. Some swathed pea crops are sprouting and producers are reporting that some pulse crops are coming off tough. At this time, four per cent of forage crops have not been cut or baled due to excess moisture or drought.

Producers are desiccating pulses, cleaning corrals and swathing and combining crops.

Southwestern Saskatchewan (Crop Districts 3ASW, 3AN, 3B and 4)

The region received very little precipitation this past week, helping to progress harvest. The Stewart Valley and Consul areas received 8 mm of rainfall, the Mankota and Kincaid areas 5 mm, the Tyner and Frontier areas 4 mm, the Admiral area 3 mm, the Ponteix and Eyebrow areas 2 mm and the Glenbain area 1 mm. Since April 1, rainfall in the region has ranged from 197 mm (Assiniboia area) to 454 mm (Cadillac area).

Twenty-seven per cent of the 2012 has now been harvested, up from 13 per cent last week. An additional 18 per cent is swathed or ready to straight cut. Seventy-seven per cent of the winter wheat, 81 per cent of the fall rye, nine per cent of the spring wheat, five per cent of the durum, 26 per cent of the barley, 60 per cent of the lentils, 75 per cent of the peas, two per cent of the chickpeas, 25 per cent of the canola and 36 per cent of the mustard have been combined. Forty-four per cent of the canola, 37 per cent of the mustard, 29 per cent of the barley, 17 per cent of the spring wheat and 12 per cent of the durum is swathed or ready to straight cut.

Average yields vary across the region depending on excess moisture in the spring, the high temperatures of July and August, and disease and insect damage. Average yields for the region are predicted as follows: winter wheat 45 bu/ac, spring wheat 31 bu/ac, durum 34 bu/ac, oats 49 bu/ac, barley 48 bu/ac, fall rye 38 bu/ac, flax 17 bu/ac, canola 27 bu/ac, mustard 960 lb./ac, lentils 1408 lb./ac, peas 32 bu/ac, canaryseed 1080 lb./ac and chickpeas 1278 lb./ac.

Moisture conditions in the region continue to worsen due to the high temperatures and lack of moisture over the past month. Topsoil moisture on cropland is rated as 34 per cent adequate, 49 per cent short and 17 per cent very short. Hay land and pasture topsoil moisture conditions are 23 per cent adequate, 45 per cent short and 32 per cent very short. In Crop District 4A, 84 per cent of the cropland, hay land and pasture is very short of topsoil moisture. This has increased from 68 per cent last week. Rainfall is needed in the majority of the region to help recharge the soil.

Much of the crop damage this past week is attributed to drought. The remaining damage is attributed to wind, insects and disease. There have been many reports of grasshoppers damaging crops and large flea beetle populations in canola. Many crops have been damaged by disease and drought and many areas are reporting disappointing yields. However, some areas that received timely rainfalls are reporting slightly-above-average yields. Strong winds last week shelled some canola and lentils fields and many swaths have been blown around. A light frost was received in some areas last Wednesday night but damage to crops has been minimal. Many crops that were harvested during high temperatures are now being aerated. At this time, two per cent of forage crops have not been cut or baled due to excess moisture or drought.

Producers are busy swathing and combining crops, hauling bales and preparing bins.

East-Central Saskatchewan (Crop Districts 5 and 6a)

Small amounts of rain this past week have delayed field operations in some areas of the region. Rainfall ranged from trace amounts to 26 mm in the Kelvington area. The Langenburg area received 14 mm of rain, the Dysart area 3 mm, the Ituna and Leroy areas 12 mm, the Rama and Kuroki areas 6 mm, the Wynyard area 2 mm, the Semans area 9 mm, the Allan and Bethune areas 1 mm and the Meacham area 4 mm. Since April 1, cumulative rainfall in the region has ranged from 244 mm (Quill Lake area) to 493 mm (Jedburgh area).

Harvest continues in the region, with four per cent of the 2012 crop combined and an additional 20 per cent swathed or ready to straight combine. Last week, only one per cent of the crop was combined and six per cent swathed or ready to straight combine. Seventy-seven per cent of the winter wheat, 89 per cent of the fall rye, one per cent of the spring wheat, five per cent of the durum, three per cent of the barley, 15 per cent of the lentils, 38 per cent of the peas, five per cent of the chickpeas and one per cent of the canola have been combined. Thirty-three per cent of the canola and six per cent of the mustard has been swathed.

Yields vary throughout the region and many crops are yielding less than what had been expected due to excess moisture, disease and high temperatures during flowering. Estimated yields of some uncombined crops have decreased in the past few weeks. Average yields for the region are predicted as follows: winter wheat 43 bu/ac, spring wheat 34 bu/ac, durum 34 bu/ac, oats 74 bu/ac, barley 55 bu/ac, fall rye 38 bu/ac, flax 22 bu/ac, canola 27 bu/ac, mustard 1146 lb./ac, lentils 1417 lb./ac, peas 33 bu/ac, canaryseed 772 lb./ac and chickpeas 1200 lb./ac.

Cropland topsoil moisture is rated as six per cent surplus, 81 per cent adequate and 13 per cent short. Hay land and pasture topsoil moisture is rated as six per cent surplus, 74 per cent adequate and 20 per cent short. Crop District 5A is reporting that 18 per cent of cropland and 16 per cent of hay land and pasture have surplus moisture.

Crop damage this week is attributed to disease, insects, hail and drought. Aster yellows, sclerotinia and bertha armyworms will affect canola yields in much of the region. Cereal yields will be less than expected due to lodging, aster yellows and sawfly damage. There have been many reports of shrunken kernels and ergot, and there are indications that quality may be an issue this year. Weed populations have increased in fields that had excess moisture and where crops have been drowned out. A light frost was received last Wednesday morning in some areas but damage is minimal. At this time, five per cent of forage crops will not be cut or baled due to excess moisture or drought; however, a second cut will be possible on some hay fields that did not receive excess moisture.

Producers are busy hauling bales, swathing and combining.

West-Central Saskatchewan (Crop Districts 6B and 7)

Varying amounts of rainfall were received in the region, delaying some field operations. The Battleford area received 20 mm, the Rosthern area 14 mm, the Arelee area 4 mm, the Conquest area 2 mm, the Rosetown area 6 mm, the Kindersley area 3 mm, the Kerrobert area 10 mm, the Landis area 8 mm and the Macklin area 17 mm. Since April 1, cumulative precipitation in the region has ranged from 310 mm (Netherhill area) to 523 mm (Outlook area). The Outlook area has the third highest cumulative rainfall total in the province since April 1.

Harvest is progressing and producers now have two per cent of the 2012 crop combined, up from less than one per cent last week. Fifteen per cent of the crop is swathed or ready to straight combine. Forty-three per cent of the winter wheat, 71 per cent of the fall rye and nine per cent of the peas have been combined. One per cent of the spring wheat, barley, lentils, canola and mustard have been combined. Twenty-eight per cent of the canola and seven per cent of the mustard is swathed. Warm and dry weather is needed to advance the maturity of many later-seeded crops.
Many crop reporters are indicating that it is difficult to assess yields at this time on uncombined crops, due to variable crop staging, flooded areas and disease. Many combined pea crops had disappointing yields due to disease and excess moisture. Average yields for the region are predicted as follows: winter wheat 41 bu/ac, spring wheat 36 bu/ac, durum 38 bu/ac, oats 68 bu/ac, barley 56 bu/ac, fall rye 30 bu/ac, flax 22 bu/ac, canola 30 bu/ac, mustard 1235 lb./ac, sunflowers 1150 lb./ac, lentils 1174 lb./ac, peas 35 bu/ac, canaryseed 882 lb./ac and chickpeas 1000 lb./ac.

Topsoil moisture conditions are rated as five per cent surplus, 84 per cent adequate and 11 per cent short on cropland. Hay land and pasture topsoil moisture conditions are rated as four per cent surplus, 77 per cent adequate and 19 per cent short.

Insects and disease caused the majority of crop damage this week. Aster yellows, sclerotinia and bertha armyworms are expected to reduce canola yields in much of the region. Pulse and cereal crops that have lodged are causing harvest issues. There have been reports of wildlife damage to many crops and some producers are still spraying for bertha armyworms in later-seeded canola. Crops are advancing quickly and additional warm weather is needed to facilitate harvest operations. At this time, two per cent of forage crops will not be cut or baled due to excess moisture.

Producers are busy swathing canola, desiccating pulses, cleaning bins and getting ready to combine.

Northeastern Saskatchewan (Crop Districts 8 and 9AE)

The region continues to receive significant amounts of rain that is delaying field operations. The Porcupine Plain area received 45 mm of rain this past week, the Arborfield and Alvena areas 13 mm, the Codette area 26 mm, the Nipawin area 31 mm, the Garrick area 30 mm, the Prince Albert area 22 mm, the Bruno area 14 mm, the Melfort area 36 mm and the Birch Hills area 24 mm. The Alvena area has recorded the least amount of rainfall since April 1 (360 mm), while the Tisdale area has received the greatest cumulative rainfall in both the region and the province (605 mm).

Just over one per cent of the 2012 crop has been combined and 11 per cent is swathed or ready to straight combine. Forty-nine per cent of the winter wheat, 81 per cent of the fall rye, two per cent of the barley, one per cent of the lentils and 15 per cent of the peas have been combined. An additional 26 per cent of the peas and 28 per cent of the canola is swathed or ready to straight combine.

Many crop reporters are indicating that it is difficult to assess yields at this time on uncombined crops, due to variable crop staging, flooded areas and disease. Average yields for the region are predicted as follows: winter wheat 40 bu/ac, spring wheat 38 bu/ac, durum 38 bu/ac, oats 81 bu/ac, barley 61 bu/ac, fall rye 43 bu/ac, flax 21 bu/ac, canola 30 bu/ac, mustard 900 lb./ac, lentils 1398 lb./ac, peas 33 bu/ac, canaryseed 1113 lb./ac and chickpeas 1200 lb./ac.

Cropland topsoil moisture is rated as 24 per cent surplus and 76 per cent adequate. Hay land and pasture topsoil moisture is rated as 12 per cent surplus and 88 per cent adequate. Crop districts 8A and 9AE are reporting 25 per cent of cropland has surplus moisture. Eighteen per cent of hay land and pasture in CD 8B has surplus moisture.

Insects, disease and localized flooding are causing most of the crop damage this week. Many fields in the region have flooded areas and cannot support equipment. Water that lies in some fields and across access roads will cause harvest challenges if warm and dry weather does not arrive soon. Spraying for bertha armyworms and diamondback moths continues in some parts of the region. Aster yellows, sclerotinia and insects will cause yield loss in many canola yields. Crops are ripening quickly and the bulk of harvest is expected to begin soon if the fields dry up. At this time, 11 per cent of forage crops will not be cut or baled due to excess moisture; however, many producers will be getting a second cut of hay.

Producers are busy hauling bales, desiccating pulses and getting ready for harvest.

Northwestern Saskatchewan (Crop Districts 9AW and 9B)

Much of the region received rainfall this past week that has delayed field operations. The Hafford area received 10 mm of rain, the Speers area 14 mm, the Duck Lake area 18 mm, the North Battleford area 20 mm, the Glaslyn area 17 mm, the Turtleford area 5 mm, the Frenchman Butte area 4 mm and the Dorintosh area 3 mm. Since April 1, rainfall in the region has varied from 236 mm (Dorintosh area) to 437 mm (Radisson area).

Harvest is just beginning in some parts of the region. Just over one per cent of the 2012 crop is combined and 11 per cent is swathed or ready for straight combining. Twelve per cent of the peas have been combined and an additional 26 per cent is swathed or ready to straight cut. Ninety-nine per cent of the winter wheat, 85 per cent of the fall rye and six per cent of the spring wheat is swathed or ready to straight cut. Most harvest operations will begin in the coming weeks.

Many crop reporters are indicating that it is difficult to assess yields at this time on uncombined crops, due to variable crop staging, flooded areas, insects and disease. Average yields for the region are predicted as follows: winter wheat 60 bu/ac, spring wheat 41 bu/ac, oats 91 bu/ac, barley 70 bu/ac, fall rye 39 bu/ac, canola 33 bu/ac, lentils 600 lb./ac and peas 36 bu/ac.

Topsoil moisture conditions for cropland are three per cent surplus, 93 per cent adequate and four per cent short. Pasture and hay land conditions are one per cent surplus, 89 per cent adequate and ten per cent short.

The majority of crop damage is attributed to wind, insects and disease. Spraying continues for bertha armyworms and lygus bugs in some areas of the region. Some producers have noticed wheat midge larvae in wheat heads but damage appears minimal at this time. There have been reports of wildlife and disease damage on many pea fields. Aster yellows have damaged most canola crops and yield loss will be significant in many areas of the region. Many crop yields will be less than anticipated due to disease, insects and heat stress. At this time, one per cent of forage crops will not be cut or baled due to drought.

Producers are busy finishing haying, hauling bales, desiccating pulses and getting ready for harvest.

Saskatchewan Harvest Progress August 20, 2012

Winter Wheat	% Standing	% in swath	% ready to straight combine	% combined
SE	4	4	9	83
SW*	10	1	11	77
EC	0	5	18	77
WC	11	6	40	43
NE	48	0	3	49
NW	1	0	99	0

* one per cent is rated as 'other'

Fall Rye	% Standing	% in swath	% ready to straight combine	% combined
SE	7	5	10	78
SW*	1	1	0	81
EC	0	7	4	89
WC	0	12	17	71
NE**	0	10	7	81
NW	15	3	82	0

* 17 per cent is rated as 'other'
** two per cent is rated as 'other'

Spring Wheat	% Standing	% in swath	% ready to straight combine	% combined
SE	74	6	8	12
SW	74	3	14	9
EC	91	4	4	1
WC	96	2	1	1
NE	96	1	3	0
NW	94	3	3	0

Durum	% Standing	% in swath	% ready to straight combine	% combined
SE	76	5	6	13
SW	83	3	9	5
EC	98	1	0	1
WC	97	2	1	0
NE	100	0	0	0
NW	N/A	N/A	N/A	N/A

Barley	% Standing	% in swath	% ready to straight combine	% combined
SE	72	12	6	10
SW	45	5	24	26
EC	83	8	6	3
WC	92	5	2	1
NE	84	6	8	2
NW	98	2	0	0

Lentils	% Standing	% in swath	% ready to straight combine	% combined
SE	37	13	10	40
SW	20	6	14	60
EC	60	2	23	15
WC	83	5	11	1
NE	82	7	10	1
NW	100	0	0	0

Peas	% Standing	% in swath	% ready to straight combine	% combined
SE	30	2	14	54
SW*	14	0	10	75
EC	36	7	19	38
WC	60	3	28	9
NE	59	11	15	15
NW	62	5	21	12

* one per cent is rated as 'other'

Chickpeas	% Standing	% in swath	% ready to straight combine	% combined
SE	29	66	0	5
SW	94	0	4	2
EC	60	12	23	5
WC	83	0	17	0
NE	N/A	N/A	N/A	N/A
NW	N/A	N/A	N/A	N/A

Canola	% Standing	% in swath	% ready to straight combine	% combined
SE	34	50	2	14
SW	31	42	2	25
EC	66	33	0	1
WC	71	28	0	1
NE	84	16	0	0
NW	82	18	0	0

Mustard	% Standing	% in swath	% ready to straight combine	% combined
SE	52	46	0	2
SW	27	15	22	36
EC	92	6	2	0
WC	91	7	1	1
NE	100	0	0	0
NW	N/A	N/A	N/A	N/A

Crop Districts and Rural Municipalities in Saskatchewan



Weekly Rainfall

for the week ending August 20, 2012



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2012 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Rainfall - Ministry of Agriculture, Crop Report Database
IDW interpolation (power 2.5, fixed radius 300 km)

Prepared by: Geomatics Services Date: August 22, 2012

Weekly Rainfall Summary

(in millimeters) 1 inch = 25 mm for the period August 14 to 20, 2012

Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
1A	1	Argyle	N/A	181
	2	Mount Pleasant	N/A	212
	3	Enniskillen	NIL	309
	33	Moose Creek	NIL	266
	34	Browning	N/A	230
	61 A	Antler	N/A	309
	61 B	Antler	N/A	26
	63	Moose Mountain	NIL	364.3
	64	Brock	NIL	326
	65	Tecumseh	NIL	326
1B	91	Maryfield	NIL	344
	92	Walpole	N/A	302
	95	Golden West	NIL	301
	122	Martin	N/A	415
	123	Silverwood	NIL	384
	124	Kingsley	1	279
	151 A	Rocanville	2	581
	151 B	Rocanville	NIL	483
	153	Willowdale	NIL	391
	154	Elcapo	NIL	272
	155	Wolseley	NIL	378
2A	66	Griffin	N/A	N/A
	67	Weyburn	N/A	208
	68	Brokenshell	NIL	252
	69	Norton	N/A	N/A
	97	Wellington	1	338
2B	127 A	Francis	2	362
	127 B	Francis	NIL	246.4
	129	Bratt's Lake	3	317.5
	131	Baildon	2	377
	156	Indian Head	2	447.5
	160 A	Pense	NIL	322
	160 B	Pense	0.5	329.5
	161	Moose Jaw	NIL	363
	162	Caron	4	395.7
	191	Marquis	NIL	302
3ASE	38 A	Laurier	NIL	285
	38 B	Laurier	NIL	233
	39 A	The Gap	NIL	190
	39 B	The Gap	NIL	244
3ASW	10	Happy Valley	NIL	240
	12	Poplar Valley	NIL	275
	40	Bengough	N/A	213
	42	Willow Bunch	N/A	296
	43	Old Post	3	299
	44	Waverley	N/A	262
	70	Key West	N/A	160
	71	Excel	NIL	197
	73 A	Stonehenge	NIL	215.3
	73 B	Stonehenge	3	265
	74	Wood River	NIL	205.6
3AN	101	Terrell	NIL	312
	102	Lake Johnston	NIL	260.1
	103	Sutton	NIL	263
	104	Gravelbourg	N/A	243.5
	132	Hillsborough	NIL	373
	134	Shamrock	NIL	248
	164	Chaplin	NIL	244
	193 A	Eyebrow	2	307
	193 B	Eyebrow	NIL	377.5
	194	Enfield	N/A	336
3BS	17	Val Marie	NIL	454
	18	Lone Tree	N/A	N/A
	45	Mankota	5	328
	75 A	Pinto Creek	5	358
	75 B	Pinto Creek	NIL	255
	76 A	Auvergne	1	309
	76 B	Auvergne	3	305
	77	Wise Creek	3	259
	78 A	Grassy Creek	3	266
	78 B	Grassy Creek	N/A	216
	78 C	Grassy Creek	5	231
	105	Glenbain	1	280
	106	Whiska Creek	3	322
	107	Lac Pelletier	N/A	262
	108	Bone Creek	4	272
3BN	138	Webb	N/A	311
	166	Excelsior	N/A	354
	167	Sask. Landing	7.8	331.8
	168	Riverside	NIL	233
	226	Victory	N/A	391
	228	Lacadena	4	287
	257	Mankota	5.5	306.7

Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
4A	19	Frontier	4	209.6
	49	White Valley	N/A	242
	51	Reno	7.6	200.2
	79	Arlington	4	219
	109	Carmichael	13	344
	110	Piapot	NIL	244
	111	Maple Creek	N/A	306
4B	139	Gull Lake	0	313
	141	Big Stick	4	320
	142	Enterprise	N/A	209
	169	Pittville	N/A	186
	231	Happyland	5	251
5A	183	Fertile Belt	NIL	456
	186	Abernethy	8	413
	211	Churchbridge	14	414
	213	Saltcoats	4	381
	214	Cana	N/A	N/A
	216	Tullymet	N/A	406
	217	Lipton	3	383
	241	Calder	4	382
	243	Wallace	NIL	326
	245 A	Garry	N/A	425
	245 B	Garry	3	493
	245 C	Garry	8	421
	246	Ituna Bon Accord	12	370
	247	Kellross	2	373
	248	Touchwood	N/A	303
5B	271	Cote	3	461
	276	Foam Lake	N/A	288.5
	277	Emerald	NIL	444
	305	Invermay	6	370
	307	Elfros	NIL	414
	308	Big Quill	2	261
	333 A	Clayton	N/A	258.8
	334	Preeceville	9	350
	336	Sasman	6	359
	337	Lakeview	N/A	332
	338	Lakeside	4	244
	366	Kelvington	26	408
6A	189	Lumsden	1	321
	190 A	Dufferin	N/A	307
	190 B	Dufferin	NIL	303
	190 C	Dufferin	NIL	274
	190 D	Dufferin	1	403.4
	219	Longlaketon	1	264
	220	Mckillop	NIL	310
	221	Sarnia	2.4	291.7
	222	Craik	2	359
	251	Big Arm	2	278
	252	Arm River	NIL	286
	279	Mount Hope	9	301
	282	McCraney	3	377
	309	Prairie Rose	N/A	283
	313	Lost River	1	300
	339	Leroy	11.6	405.1
	340	Wolverine	7	344
	341	Viscount	4	336
	343	Blucher	2	342.5
	283	Rosedale	2	314.5
6B	284	Rudy	4	523
	285	Fertile Valley	2	442
	286	Milden	NIL	396
	314	Dundurn	2	336
	344	Corman Park	2	400
	346	Perdue	N/A	454
	376 A	Eagle Creek	4	393
	376 B	Eagle Creek	9	520
	403	Rosthern	14	405
	259	Snipe Lake	2	410.1
7A	260 A	Newcombe	N/A	238
	260 B	Newcombe	N/A	310
	287	St. Andrews	6	413
	288	Pleasant Valley	7	408
	290	Kindersley	4	310
	292	Milton	N/A	410.5
	317 A	Manitou	N/A	373
	317 B	Marriott	5	404
	318	Mountain View	NIL	404
	320 A	Oakdale	3	343
	320 B	Oakdale	N/A	292
	321	Prairiedale	NIL	323

Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
7B	347 A	Biggar	NIL	395
	347 B	Biggar	NIL	358
	350 A	Manposa	10.2	463.4
	350 B	Manposa	7	351
	351	Progress	8	391
	352	Heart's Hill	NIL	426
	377	Glenside	9	479
	378 A	Rosemount	7	434
	378 B	Rosemount	8	438
	379	Reford	8	419
	381	Grass Lake	12	407
	409 A	Buffalo	20	514
	409 B	Buffalo	N/A	356.8
	410	Round Valley	1	311.2
	411	Senlac	17	437
8A	395	Porcupine	45	423
	397	Barrier Valley	N/A	253
	428	Star City	16	426
	456	Arborfield	13	501
	457	Connaught	21	605
	458	Willow Creek	26	406
	486	Moose Range	28	440
	487	Nipawin	34	528
8B	369	St. Peter	22	422
	370	Humboldt	10	422
	371	Bayne	14	459
	372	Grant	9.5	414.9
	402	Fish Creek	13	360
	429	Flett's Springs	36	446
	459	Kinistino	35	582
	460	Birch Hills	24.3	467
9AE	461	Prince Albert	22	433.5
	488	Torch River	30	399
	491	Buckland	N/A	379.9
9AW	406	Mayfield	8	437
	435	Redberry	10	428
	436	Douglas	14	332
	463	Duck Lake	18	403
	464	Leask	N/A	15
	467 A	Round Hill	20	319
	467 B	Round Hill	16	318
	494	Canwood	N/A	324
	497	Medstead	N/A	368
9B	438	Battle River	16	418
	440	Hillsdale	N/A	358.5
	442	Manitou Lake	19.8	340.3
	468 B	Meota	N/A	38
	498 A	Parkdale	17	409
	498 B	Parkdale	NIL	325
	499	Mervin	5	338.4
	501 A	Frenchman Butte	4.5	304.5
	501 B	Frenchman Butte	3	260
	501 C	Frenchman Butte	4	352
	561	Loon Lake	2	260
	588 A	Meadow Lake	4	276
	588 B	Meadow Lake	5	264
	588 C	Meadow Lake	3	236
	588 D	Meadow Lake	NIL	310.6
	622	Beaver River	N/A	275

Municipality No: A, B, C, and D - more than one reporter

These precipitation amounts represent point locations within each municipality and do not necessarily reflect the whole R. M.

Cumulative Rainfall

From: April 1, 2012
To: August 20, 2012



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2012 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Rainfall - Ministry of Agriculture, Crop Report Database
IDW interpolation (power 2.5, fixed radius 300 km)

Prepared by: Geomatics Services Date: August 22, 2012

Cropland Topsoil Moisture Conditions

August 21, 2012



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2012 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Moisture - Ministry of Agriculture, Crop Report Database
IDW interpolation (power 2.5, fixed radius 300 km)

Prepared by: Geomatics Services Date: August 22, 2012

Hay and Pasture Topsoil Moisture Conditions

August 21, 2012



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2012 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Moisture - Ministry of Agriculture, Crop Report Database
IDW interpolation (power 2.5, fixed radius 300 km)

Prepared by: Geomatics Services Date: August 22, 2012